BLACK EARTH

humiZEN™ Fuze DS

CONTAINS BENEFICIAL SUBSTANCE

Humic Acids derived from humalite	2.0%

PRODUCT INFORMATION

HumiZEN™ Alpha is a liquid solution containing humic acids. It is suitable for applications as a soil amendment and foliar spray and is compatible with a wide range of liquid fertilizers. Applications made according to directions for use may aid in the uptake of micronutrients.

DIRECTIONS FOR USE

Read label in its entirety. Shake or agitate container prior to use. Apply humiZEN™ Alpha as a stand-alone product or blended with compatible liquid fertilizers. Suitable applications include banding, injection into irrigation water, blending with transplant water, and foliar sprays. Always conduct a jar test to confirm compatibility prior to blending with any other product. Apply only as directed.

Refer to Technical Data Sheet (TDS) for specific blending/mixing capabilities for this product.

Information regarding the contents and levels of metals in this product is available at https://www.aapfco.org/metals.html

PRODUCT OF CANADA / PRODUIT DU CANADA

HPTA TEST METHOD CERTIFIED

MADE WITH CANADIAN HUMALITE™ VERIFIED SOURCE HUMIC RICH CARBON

OMRI® LISTED

RECOMMENDED APPLICATIONS

CEREALS, OILSEEDS, PULSES, FRUIT & VEGETABLE

Soil applications: Apply in band applications, in furrow, or injected into irrigation water at rates of .27–1.07 gal./ac. per application, up to 5 times per season, from pre-plant until fruit set.

Foliar applications: Apply at rates of .134-.267 gal./ac. per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from pre-plant until fruiting. **Caution** – applications made after fruit set may stain fruit.

TREE FRUITS & VINES

Soil applications: Apply broadcast or band applications, or injected into irrigation water, at rates of .27-1.07 gal./ac. per application, up to 5 times per season. Applications may be made at any point in the growing season, until and including fruit set.

Foliar applications: Apply at rates of .134-.267 gal./ac. per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from pre-plant until fruiting. **Caution** – applications made after fruit set may stain fruit.

TURF & ORNAMENTAL, GREENHOUSE CROPS (INCLUDING CANNABIS)

Transplant water: Add to transplant water at rates 1 gal./100 gal. of water.

Soil applications: Apply broadcast or band applications, or injected into irrigation water, at rates .27-1.07 gal./ac. per application, up to 5 times per season at any growth stage

Foliar applications: Apply at rates of .134-.267 gal./ac. per application, diluted in water at maximum 10% concentration, up to 5 times annually at any time from pre-plant until fruit set. **Caution** – applications made after fruit set may stain fruit (or other harvestable parts of the plant).

GUARANTEED BY

WESTMET GROUP CANADA LTD

5555 Calgary Trail NW, Suite 1270, Edmonton, AB, T6H 5P9, Canada

CONDITIONS OF SALE & LIMITATION OF WARRANTY & LIABILITY

The manufacturer guarantees only that the composition of this product conforms to the description on the label and that the product is suited to the stated purposes when used according to the Direction for Use. No individuals or companies are authorized to make any further warranty, expressed or implied, concerning the selection, use, handling, or storage of this product. The manufacturers' liability will be limited to the refund of purchase price or, at manufacturers' discretion, replacement of product.

NET WEIGHT	☐ 2,315 LB (1,050 KG)	☐ 1,158 LB. (525 KG)	☐ 262 LB. (119 KG)	☐ 23 LB. (10.5 KG)
NET VOLUME	☐ 265 GAL. (1,000 L)	☐ 132 GAL. (500 L)	☐ 30 GAL. (113 L)	☐ 2.5 GAL. (10 L)

DENSITY **8.43 LB./GALLON @ 68° FAHRENHEIT**

MADE IN ALBERTA, CANADA

For distribution in the states of California & Oregon only.